



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

respect to acute intestinal diseases, which caused 33 deaths. There was, however, a notable decrease with regard to acute diseases of the respiratory organs, which claimed 60 victims. Furthermore, there were registered 4 deaths from influenza, 8 deaths from measles, 6 deaths from diphtheria, 4 deaths from scarlet fever, 76 deaths from phthisis pulmonalis, and 41 deaths from cancer. Finally, 11 persons died by violence.

Typhoid fever in Germany—Precautions to be observed.

[Extract from the "Gesundheitsbüchlein," issued by the Imperial Health Department of Germany.]

Typhoid fever.—Thanks to the improvements in the public sanitation in many large German cities in which formerly numerous cases of illness and death and even widespread epidemics were caused by typhoid fever, this disease has become rarer. It prevails, however, largely in the country and in many cities. In the twenty years from 1877 to 1896 49,948 persons died from typhoid fever in the towns of Germany which have population of not less than 15,000, being a yearly average of 2,497 deaths for the whole Empire; still, since 1877, there has been a steady decline in the number of deaths from typhoid fever. In the five years from 1877 to 1891 the average was only 2,269 annually, although there has been a great increase in the urban population during the same period; from 1892 to 1896 the average was 1,666, and in the year 1897 there were only 1,552 deaths from typhoid fever.

The typhoid germs leave the body of the patient with the excreta and urine, and easily pass, even with careful watching, to his linen and bedding. Sometimes the excretions occur involuntarily.

From his linen the infectious germs may spread to his hands, and next to all objects touched by him, such as clothes, food, and utensils, and may thus carry infection to relatives, nurses, physicians, and other persons who do not carefully observe the regulations necessary in the intercourse with the patient. The patient's linen must be disinfected as soon as possible after use, and the sick room and its furniture after the close of the illness. The excreta of the patient should never be discharged or removed without previous disinfection. The nonobservance of this latter regulation is a frequent cause of epidemics of typhoid fever.

HONDURAS.

Epidemic of anthrax in department of Intibuca.

The following is received from Consul Moe at Tegucigalpa, under date of December 22:

Late in November a disease broke out among the inhabitants of the department of Intibuca, Honduras, and became general in the town of La Esperanza. At first the disease was variously diagnosed, and as little was done in the matter of confining its spread or alleviating the suffering of those persons afflicted many became violently ill, and a large number died through nontreatment or mistreatment.

The attention of the Government was finally called to the matter, and a sanitary and hygienic commission, composed of two physicians and surgeons of high repute in Honduras, was appointed by the Government to investigate and report on the nature of the epidemic and